

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 410 498 A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90201706.0

(22) Date of filing: 27.06.90

(51) Int. Cl.⁵: **C12N 15/56**, C12N 9/28,
C12N 9/30, C12N 9/32,
C12N 15/75, D06L 1/06,
C08B 30/12, //(C12N9/28,
C12R1:07,1:10)

(30) Priority: 29.06.89 EP 89201735

(43) Date of publication of application:
30.01.91 Bulletin 91/05

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
06.11.91 Bulletin 91/45

(71) Applicant: **GIST-BROCADES N.V.**
Wateringseweg 1 P.O. Box 1
NL-2600 MA Delft(NL)

Applicant: **PLANT GENETIC SYSTEMS, N.V.**
Kolonel Bourgstraat 106 Bus 1
B-1040 Brussel(BE)

(72) Inventor: **Quax, Wilhelmus Johannes**
Jan van Galenlaan 8
NL-2253 VB Voorschoten(NL)
Inventor: **Laroche, Yves**
Rue Bemel 115
B-1150 Brussels(BE)
Inventor: **Vollebregt, Adrianus Wilhelmus**
Hermanus
Bereklaauw 13
NL-2671 WZ Naaldwijk(NL)
Inventor: **Stanssens, Patrick**
De Gieylaan 13
St. Denijs Westrem(BE)
Inventor: **Lauwereys, Marc**
Wilgenstratt 2
B-2950 Haaltert(BE)

(74) Representative: **Huygens, Arthur Victor, Dr. et al**
c/o Gist-Brocades N.V. Patent & Trademarks
Department Wateringseweg 1 PO Box 1
NL-2600 MA Delft(NL)

(54) Mutant microbial α -amylases with increased thermal, acid and/or alkaline stability.

(57) Thermostable and acid stable α -amylases are provided as expression products of genetically engineered α -amylase genes isolated from microorganisms, preferably belonging to the class of Bacilli. Both chemical and enzymatic mutagenesis methods are e.g. the bisulphite method and enzymatic misin-

corporation on gapped heteroduplex DNA. The mutant α -amylases have superior properties, e.g. improved thermostability over a broad pH range, for industrial application in starch processing and textile desizing.

EP 0 410 498 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 20 1706

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	PROTEIN ENGINEERING, vol. 1, no. 3, June 1987, page 241, Eynsham, Oxford, GB; T. GLUMOFF et al.: "A truncated, thermostable Bacillus stearo-thermophilus alpha-amylase" * Abstract no. 60 * ---	1,2,5,6 ,9-12, 15-21	C 12 N 15/56 C 12 N 9/28 C 12 N 9/30 C 12 N 9/32 C 12 N 15/75 D 06 L 1/06 C 08 B 30/12 //
X,D	JOURNAL OF BACTERIOLOGY, vol. 166, no. 2, May 1986, pages 635-643, American Society for Microbiology, Washington, DC, US; G.L. GRAY et al.: "Structural genes encoding the thermophilic alpha-amylases of Bacillus stearothermophilus and Bacillus licheniformis" * Figure 5 * ---	1,2,5,6 ,9-12, 15-21	(C 12 N 9/28 C 12 R 1:07 C 12 R 1:10)
A,D	EP-A-0 134 048 (GIST-BROCADES N.V.) ---		
X,D	EP-A-0 285 123 (SUOMEN SOKERI OY) * Page 12, lines 51-57; page 13, line 15 * ---	1,2,3,5 ,6,9-12 ,15-21	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
A,D	EP-A-0 252 666 (NOVO INDUSTRI) ---		C 12 N
A	EP-A-0 208 491 (GENENCOR INC.) * Page 1, lines 5-10 * ---		
A	WO-A-8 901 520 (CETUS CORP. et al.) ---		
P,X	PROTEIN ENGINEERING, vol. 3, no. 3, January 1990, pages 181-191, Eynsham, Oxford, GB; L. HOLM et al.: "Random mutagenesis used to probe the structure and function of Bacillus stearothermophilus alpha-amylase" * Whole document * --- -/-	1-6	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 15-07-1991	Examiner VAN PUTTEN A.J.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons -/-: member of the same patent family, corresponding document	



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions, namely:

1. claims 1(part.); 2 ; 4-13(part.); 15-21(part.)

A mutant alpha-amylase, that is an expression product of a mutated DNA sequence encoding an alpha-amylase, said mutant alpha-amylase exhibits improved thermostability for application in the degradation of starch and/or textile desizing.

2. claims 1(part.); 3; 4-13(part.); 15-21(part.)

A mutant alpha-amylase, that is an expression product of a mutated DNA sequence encoding an alpha-amylase, said mutant alpha-amylase exhibits improved stability at a pH below 6.5 and/or above 7.5 for application in the degradation of starch and/or textile desizing.

3. claim 14

A Bacillus/E.coli shuttle vector, wherein the expression of the cloned gene in E.coli is made impossible by physical separation of the regulatory sequences from the structural gene and wherein the expression of the cloned gene in Bacillus can be restored by digestion with a single restriction enzyme and subsequent recircularization.

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☒ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims: points 1. and 2..
- ☐ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
P,X	JOURNAL OF BIOCHEMISTRY, vol. 107, no. 2, February 1990, pages 267-272, Tokyo, JP; M. VIHINEN et al.: "Site-directed mutagenesis of a thermostable alpha-amylase from bacillus stearothermophilus: Putative role of three conserved residues" * Whole document *	1-6	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
The present search report has been drawn up for all claims.			
Place of search THE HAGUE		Date of completion of the search 15-07-1991	Examiner VAN PUTTEN A.J.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 410 498 A2**

12

EUROPEAN PATENT APPLICATION

21 Application number: 90201706.0

22 Date of filing: 27.06.90

51 Int. Cl.⁵: **C12N 15/56**, C12N 9/28,
C12N 9/30, C12N 9/32,
C12N 15/75, D06L 1/06,
C08B 30/12, //(C12N9/28,
C12R1:07,1:10)

30 Priority: 29.06.89 EP 89201735

43 Date of publication of application:
30.01.91 Bulletin 91/05

84 Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

71 Applicant: **GIST-BROCADES N.V.**
Wateringseweg 1 P.O. Box 1
NL-2600 MA Delft(NL)

Applicant: **PLANT GENETICS SYSTEMS, N.V.**
Kolonel Bourgstraat 106 Bus 1
B-1040 Brussel(BE)

72 Inventor: **Quax, Wilhelmus Johannes**
Jan van Galenlaan 8
NL-2253 VB Voorschoten(NL)
Inventor: **Laroche, Yves**
Rue Bemel 115
B-1150 Brussels(BE)
Inventor: **Vollebregt, Adrianus Wilhelmus**
Hermanus
Bereklaauw 13
NL-2671 WZ Naaldwijk(NL)
Inventor: **Stanssens, Patrick**
De Gieyalaan 13
St. Denijs Westrem(BE)
Inventor: **Lauwereys, Marc**
Wilgenstratt 2
B-2950 Haaltert(BE)

74 Representative: **Huygens, Arthur Victor, Dr. et**
al
c/o Gist-Brocades N.V. Patent & Trademarks
Department Wateringseweg 1 PO Box 1x 1
NL-2600 MA Delft(NL)

54 Mutant microbial alpha-amylases with increased thermal, acid and/or alkaline stability.

57 Thermostable and acid stable α -amylases are provided as expression products of genetically engineered α -amylase genes isolated from microorganisms, preferably belonging to the class of Bacilli. Both chemical and enzymatic mutagenesis methods are e.g. the bisulphite method and enzymatic misincorporation on gapped heteroduplex DNA. The mutant α -amylases have superior properties, e.g. improved thermostability over a broad pH range, for industrial application in starch processing and textile desizing.

EP 0 410 498 A2

MUTANT MICROBIAL α -AMYLASES WITH INCREASED THERMAL, ACID AND/OR ALKALINE STABILITYINTRODUCTIONTechnical Field

The present invention relates to the field of genetic engineering and provides new DNA molecules comprising DNA sequences coding for enzymes with α -amylase activity. Specifically, mutant microbial α -amylases are disclosed having improved characteristics for use in the degradation of starch, in the desizing of textile and in other industrial processes. The disclosed α -amylases show increased thermal, acid and alkaline stability which makes them ideally suited for performing their activity under process conditions which could hitherto not be used.

Background of the invention

Starch consists of a mixture of amylose (15-30% w/w) and amylopectin (70-85% w/w). Amylose consists of linear chains of α -1,4-linked glucose units having a molecular weight (MW) from about 60,000 to about 800,000. Amylopectin is a branched polymer containing α -1,6 branch points every 24-30 glucose units, its MW may be as high as 100 million.

Sugars from starch, in the form of concentrated dextrose syrups, are currently produced by an enzyme catalyzed process involving: (1) liquefaction (or thinning) of solid starch with an α -amylase into dextrins having an average degree of polymerization of about 7-10, and (2) saccharification of the resulting liquefied starch (i.e. starch hydrolysate) with amyloglucosidase (also called glucoamylase or AG). The resulting syrup has a high glucose content. Much of the glucose syrup which is commercially produced is subsequently enzymatically isomerized to a dextrose/fructose mixture known as isosyrup.

α -Amylase (EC 3.2.1.1) hydrolyzes starch, glycogen and related polysaccharides by cleaving internal α -1,4-glucosidic bonds at random. This enzyme has a number of important commercial applications in, for example the sugar, brewing, alcohol and textile industry. α -Amylases are isolated from a wide variety of bacterial, fungal, plant and animal sources. The industrially most important α -amylases are those isolated from Bacilli.

In the first step of the starch degradation process, starch slurry is gelatinized by heating at relatively high temperature (up to 110°C). The gelatinized starch is liquefied and dextrinized by a thermostable α -amylase in a continuous two stage process. The major process variables are starch concentration, α -amylase dose, temperature and pH. During the liquefaction-dextrinization reaction the process variables must be maintained within narrow limits to achieve good conversion ratios, since serious filtration problems may arise otherwise. See, for example, L.E. Coker and K. Venkatasubramanian, in: Biotechnology, p. 165-171, Ed. P.N. Cheremisinoff, P.B. Quелlette, Technicom Publ. Corp. Lancaster Penn. 1985. One of the problems which frequently arises is the proper regulation of the temperature in the initial stage of the degradation process: overheating often causes denaturation of the α -amylase so that the final thinning is not sufficient. One way to avoid this is the use of more thermostable α -amylases.

To that end it has been proposed to add calcium ions or an amphiphile (see e.g. EP-A-0189838), but this solution appeared to be unsatisfactory.

There is, therefore, still substantial interest to provide α -amylases with increased thermostability.

Relevant Literature

EP-A-057976 describes the isolation of a thermostable α -amylase coding gene from *B. stearothermophilus* the gene is cloned into a plasmid containing either a *Bacillus* or an *E. coli* origin of replication. The so obtained chimeric plasmid is used for producing α -amylase. The α -amylase gene was isolated and used without any further modification.

EP-A-0134048 describes a method for increased commercial production inter alia of α -amylase, by cloning and expression of one or more α -amylase genes in industrial Bacillus strains.

EP-A-252666 describes a chimeric α -amylase with the general formula Q-R-L in which Q is a N-terminal polypeptide of 55 to 60 amino acid residues which is at least 75 percent homologous to the 37 N-terminal residues of the B. amyloliquefaciens α -amylase, R is a given polypeptide and L is a C-terminal polypeptide of 390 to 400 amino acid residues which is at least 75 percent homologous to the 395 C-terminal residues of B. licheniformis α -amylase.

Gray et al. (J. Bacteriol., 1986, 166, 635) describe chimeric α -amylases formed of the NH₂-terminal portion of B. stearothermophilus α -amylase and the COOH-terminal portion of B. licheniformis α -amylase. Most of the hybrid enzyme molecules were shown to be less stable than the parent wild-type enzymes. Furthermore none of the hybrid molecules was shown to possess improved stability properties.

None of the references cited above describes the use of single amino acid replacements to obtain novel α -amylases.

EP-A-0285123 discloses a method for complete mutagenesis of nucleic acid sequences. As an example mutagenesis of the B. stearothermophilus α -amylase is described. Although there is a suggestion that this method can be used to obtain B. stearothermophilus α -amylase mutants with improved stability no examples are given.

SUMMARY OF THE INVENTION

The present invention provides mutant α -amylases and ways of obtaining such mutants. Said mutant α -amylases are characterized in that they differ in at least one amino acid from the wild-type enzyme. Furthermore, DNAs encoding these mutants, vectors containing these DNAs in expressionable form and host cells containing these vectors are provided.

In one aspect of the invention random mutagenesis on cloned α -amylase genes is disclosed. The mutated genes are expressed in a suitable host organism using a suitable vector system.

In another aspect of the invention screening methods for mutant α -amylases are described and applied. Said methods yield more thermostable and more acid stable α -amylases. Furthermore, this method is used with a slight modification to obtain more alkaline stable α -amylases. The expression products of the clones so detected are isolated and purified.

In yet another aspect of the invention α -amylases are provided with increased thermostability, these mutant α -amylases reduce filtration problems under application conditions of starch degradation.

In a further aspect of the invention α -amylases are provided with increased acid stability, these reduce the formation of unfavourable by-products, such as maltulose, at the same time they decrease the amount of acid to be added before the reaction with amyloglucosidase. The new α -amylases possess preferably both improved properties with respect to thermostability and acid stability or with respect to both thermostability and alkaline stability.

In another aspect of the invention the mutant proteins are shown to have a better performance under application conditions of starch liquefaction. The alkaline stability is especially useful for application in textile desizing.

These aspects will be further described in the detailed description and in the examples hereinafter.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Nucleotide sequence of pMa5-8

Stanssens et al., 1987, EMBO Laboratory Course Martinsried, July 1987. For description of the different elements see text.

Figure 2: Nucleotide sequence of plasmid pPROM SPO2 insert

Construction of this vector has been described in EP-A-0224294. The α -amylase amino acid sequence is depicted below the triplets. Numbering starts from the first amino acid of the mature protein (Kuhn et al., 1982, J. Bacteriol., 149, 372). The SPO2 promoter insert runs from position 61 to 344.

Figure 3: Nucleotide sequence of pMaTLia6

This vector was constructed from pMa5-8, the insert of pPROM SPO2 and a synthetic DNA fragment encoding the TAC promoter. The TAC promoter DNA fragment runs from position 3757 to position 3859.

The α -amylase amino acid sequence is depicted below the triplets.

Figure 4 :Restriction map of pMaTLia6

The following unique restriction enzyme sites are available for gap construction in the α -amylase gene: Bam HI, Spe I, Sac II, Kpn I, Cla I, Nar I, Sal I, Tht 111I, Xma III and Bst EII. Sequencing primers for all possible gaps have been synthesized in order to enable easy determination of mutations. Plasmid pMcTLia6 is identical with pMaTLia6 except for the presence of an amber codon in the ampicillin gene (removes Sca I site) and the absence of an amber codon in the chloramphenicol gene (associated with the presence of a Pvu II site).

Figure 5: Outline of Bacillus/E. coli shuttle vector pBMa/c

The (left) pMa/c section enables convenient mutagenesis in E. coli. The (right) Bacillus subtilis cassette contains the α -amylase gene (or any other Bacillus gene) plus a minimal replicon for propagation in B. subtilis. After successful mutagenesis in E. coli the B. subtilis cassette can be circularized allowing the SPO2 promoter to move in front of the α -amylase gene upon transformation into Bacillus.

Figure 6: Restriction map of pBMa/c1

This vector is a specific example of the mutagenesis expression vector outlined in Figure 5.

(1) and (2): multiple cloning sites. The target gene is inserted in (2). By varying the sites at (1) and (2) convenient restriction sites for gapped duplex creation can be constructed;

FDT : transcription terminator

F1.ORI : origin of replication originating from phage F1

E. coli ORI: origin of replication from pBR322

BLA : ampicillin resistance gene

CAT : chloramphenicol resistance gene

BAC ORI : origin of replication of pUB110

KANAMYCIN : kanamycin (neomycin) resistance gene of pUB110

SPO2 : promoter of phage SPO2

Figure 7: Restriction map of pBMa/c6Lia6

The Bacillus licheniformis α -amylase gene was engineered into pBMa/c1 at multiple cloning site (2) of Figure 6. In this figure the SPO2 promoter is indicated by (2) and the E. coli ORI is represented by (4).

Figure 8: Sequence of phoA signal sequence fragment in pMa/c TPLia6

Depicted is the sequence from the Eco RI site upstream from the TAC-promoter up to the first amino acids of mature α -amylase. The phoA amino acid sequence is shown below the DNA sequence.

Figure 9: Michaelis-Menten plot for WT and 2D5 α -amylase

This plot shows the initial rate of enzyme activities vs. substrate concentration for WT and 2D5 α -amylase. Assay conditions are described in Example 8.

Figure 10: Thermoinactivation of WT and D7 α -amylase

This plot shows the half life time of both WT and D7 α -amylase as a function of the Ca^{2+} concentration at pH 5.5 and 90.5 °C.

Figure 11: Thermoinactivation of WT and D7 α -amylase

As in Figure 10 except for the pH which is 7.0.

Figure 12: Thermoinactivation of WT and 2D5 α -amylase

This plot shows half life times of both WT and 2D5 α -amylase as a function of Ca^{2+} concentration at pH 7.0 and 95 °C.

Figure 13: Thermoinactivation of WT and D7 α -amylase as a function of pH

Figure 14: Thermoinactivation of WT and 2D5 α -amylase as a function of pH

Figure 15: DE vs final pH measured after liquefaction at 110 °C

DETAILED DESCRIPTION OF THE INVENTION

By the term "exhibits improved properties" as used in connection with "mutant α -amylase" in the present description we mean α -amylases which have a higher enzymatic activity or a longer half-life time under the application conditions of starch liquefaction, textile desizing and other industrial processes.

With "improved thermostability" we mean that the mutant enzyme retains its activity at a higher process temperature, or that it performs longer at the same temperature than the wild-type enzyme from which it originates.

With "improved acid (or alkaline) stability" we mean that the mutant enzyme performs better at lower (or higher) pH values than the wild-type enzyme from which it was derived.

It is to be understood that the improved properties are caused by the replacement of one or more

amino acids.

Chromosomal DNA may be isolated from an α -amylase containing microorganism. Preferably a microorganism is used belonging to the genus *Bacillus*, more preferably *B. licheniformis*, still more preferably *B. licheniformis* T5 is used (see EP-A-134048). The chromosomal DNA is digested with a suitable restriction enzyme and cloned into a vector. A number of possible ways of selection can be used e.g. hybridization, immunological detection and detection of enzymatic activity. The choice of the vector used for cloning the digested chromosomal DNA will depend on the selection method available. If hybridization is used no special precautions are needed. However, if detection is immunological or based on enzymatic activity the vector will have to contain the proper expression signals. The actual detection of clones containing α -amylase was performed on starch containing agar plates. After growth and incubation with I_2 vapor halos are detected around positive clones. As a next step the sequence of the gene is determined. The derived amino acid sequence is used for comparison with other known α -amylase sequences to give a first impression of important amino acids (e.g. active-site, Ca^{2+} binding, possible S-S bridges). A better indication is obtained when the 3D-structure is determined. Since this is very laborious oftentimes another approach is used. In the absence of a 3D-structure prediction programs for determining the secondary structural elements (e.g. α -helix, β -sheet) are successfully used eventually the tertiary structural elements e.g. β -barrel are determined. For a review see Janin, J. and Wodack, S.J., Prog. Biophys. molec. Biol. 1983, 42, 21-78.

Valuable amino acid replacements can be envisioned. The stability of a protein structure is determined by the net difference in free energy between the folded and unfolded conformations of the protein. Since the proline residue is restricted to fewer conformations than the other amino acids the configurational entropy of unfolding a protein is decreased (and stability thereby increased) when an amino acid is replaced with proline. Another useful substitution is the glycine to alanine replacement. Residues such as threonine, valine and isoleucine with branched β -carbons restrict the backbone conformation more than non-branched residues.

Since a part of the thermostability of certain proteins is due to salt bridges it may be advantageous to introduce lysine and arginine residues (Tomozic S.J. and Klibanov A.M., J. Biol. Chem., 1988, 263 3092-3096). Moreover replacement of lysine by arginine residues may improve the stability of salt bridges since arginine is able to form an additional H-bond. For a review see Wigby, D.B. et al. Biochem. Biophys. Res. Comm. 1987, 149, 927-929. Deamidation of asparagine and glutamine is mentioned to cause a serious disruption of the enzyme structure, replacement with non-amide residues may avoid this disruption. Amino acid replacements are best made by mutagenesis at the DNA level.

In principle mutagenesis experiments can be performed immediately on isolated clones. However, the insert is preferably cloned in a mutagenesis/expression vector. Random mutagenesis is possible and so is site-directed mutagenesis. In view of the huge amount of mutated clones of the former method, and since no 3D-structure of α -amylase is known to make possible an educated guess for site-directed mutagenesis we decided to perform "random" mutagenesis in specific regions.

The following is a possible approach for practising the present invention.

First the gene is modified by the introduction of "silent" restriction sites. Introduction of non-silent restriction sites is also possible. This makes possible the deletion of specific regions of the gene. Secondly the gene is cloned in a phasmid. This combination of a phage and a plasmid makes easy the production of single stranded DNA. Other ways of obtaining single stranded DNA are also possible. By hybridizing melted double-stranded vector (plus insert) DNA with a vector/insert combination containing a gap in the insert, gapped heteroduplex DNA was obtained (for a detailed description see Morinaga, Y et al. 1984, Biotechnology, 2, 636).

The gap is used for chemical or enzymatic mutagenesis. Preferably we used the bisulphite method (Folk and Hofstetter, Cell, 1983, 33, 585) and an enzymatic misincorporation method are used (modified version of Lehtovaara et al., Prot. Eng., 1988, 2, 63). These methods can be applied in such a way that every single nucleotide in the gap is replaced by all three other nucleotides (saturation mutagenesis). The latter method can be applied in several ways. In one of them a synthetic primer is hybridized to the gap. Subsequently an extension reaction is performed in which the deoxynucleotide complementary to the first deoxynucleotide 3' from the primer is missing. In principle all three of the other deoxynucleotides can thus be incorporated. This can be achieved either by using a mix of three deoxynucleotides or by using three separate reactions each containing only one deoxynucleotide. Another way of applying the method yields random clones. Here, four separate reactions are set up each of them containing one limiting deoxynucleotide. This gives second strands that stop before every single nucleotide. The subsequent steps can be performed as described above. Both the bisulphite and the enzymatic mutagenesis method were employed to obtain mutants.

For testing the enzymatic properties it may be convenient to express the cloned genes in the same host as that used during mutagenesis experiments. In principle this can be any host cell provided that suitable mutagenesis/expression vector systems for these cells are available. For the most part *E. coli* is very convenient to work with, for example *E. coli* WK6. After growth of the colonies in microtiterplates samples from the wells of these plates are spotted on agar plates supplemented with starch and buffered at different pH values. Positive clones can be detected by halo formation. Screening with appropriate buffers can be used to select for thermostability, acid stability, alkaline stability, saline stability or any other stability that can be screened for.

Suitable host strains for production of mutant α -amylases include transformable microorganisms in which the expression of α -amylase can be achieved. Specifically host strains of the same species or genus from which the α -amylase is derived, are suited, such as a *Bacillus* strain. Preferably an α -amylase negative *Bacillus* strain is used more preferably an α -amylase and protease negative *Bacillus* strain.

For example *B. licheniformis* T9 has been used to produce high amounts of mutant α -amylases.

Preferably, the α -amylases being produced are secreted into the culture medium (during fermentation), which facilitates their recovery. Any suitable signal sequence can be used to achieve secretion.

The expressed α -amylase is secreted from the cells and can be subsequently purified by any suitable method. Gelfiltration and Mono Q chromatography are examples of such methods. The isolated α -amylase was tested for thermoinactivation at different Ca^{2+} concentrations (0.5 - 15 mM) and over a wide pH range (5.5 - 8.0). Tests were also performed under application conditions. Specifically mutant α -amylase was tested under conditions of starch liquefaction at pH 5.5 and 5.25. Furthermore, applications for textile desizing have been tested.

The properties of some of the mutants that are screened will be better suited under the desired performance conditions.

The present invention discloses α -amylases with increased thermostability, improved acid stability and improved alkaline stability. Generally the number of amino acid replacements is not important as long as the activity of the mutated protein is the same or better than that of the wild-type enzyme. Mutant α -amylases differ in at least one amino acid from the wild-type enzyme, preferably the mutants differ in from 1 to 10 amino acids. Specific mutants with improved properties include mutant α -amylases containing one or more amino acid replacements at the following positions 111, 133 and 149 (numbering is in accordance with the *B. licheniformis* α -amylase). Among the preferable amino and replacements are Ala-111-Thr, His-133-Tyr and Thr-149-Ile.

Such mutant enzymes show an improved performance at pH values below 6.5 and/or above 7.5. The performance is also increased at high temperatures leading to an increased half-life-time at for example temperatures of up to 110 °C.

Many of the available α -amylase products are obtained from bacterial sources, in particular *Bacilli*, e.g. *B. subtilis*, *B. licheniformis*, *B. stearothermophilus*, *B. coagulans* and *B. amyloliquefaciens*. These enzymes show a high degree of homology and similarity (Yuuki et al., J. Biochem., 1985, 98, 1147; Nakajima et al., Appl. Microbiol. Biotechnol., 1986, 23, 355). Therefore knowledge of favourable mutations obtained from one of these α -amylases can be used to improve other amylases. The present invention provides an approach for obtaining such knowledge.

Following is a description of the experimental methods used and examples to illustrate the invention. The examples are only for illustrative purpose and are therefore in no way intended to limit the scope of the invention.

EXPERIMENTAL

Materials and Methods

1. General cloning techniques

Cloning techniques were used as described in the handbooks of T. Maniatis et al., 1982, Molecular Cloning, Cold Spring Harbor Laboratory; F.M. Ausubel et al., 1987, Current Protocols in Molecular Biology, John Wiley & Sons Inc., New York; B. Perbal, 1988, A practical Guide to Molecular Cloning, 2nd edition,

John Wiley & Sons Inc., New York. These handbooks describe in detail the protocols for construction and propagation of recombinant DNA molecules, the procedures for making gene libraries, the procedures for sequencing and mutating DNA and the protocols for the enzymatic handling of DNA molecules.

5

2. Chemical mutagenesis

Cloned DNA may be treated in vitro with chemicals in order to introduce mutations in the DNA. If these mutations are directed to amino acid encoding triplet codons a mutated protein can be produced by the mutated cloned DNA. A method for chemical mutagenesis with the aid of sodium bisulfite is described by Shortle and Botstein (Methods Enzymol., 1983, 100, 457). A preferable method is described by Folk and Hofstetter (Cell, 1983, 33, 585). Other methods for mutagenesis are described by Smith, Ann. Rev. Genet., 1985, 19, 423. A particularly useful protocol is described by Ausubel et al., *ibid.*

15

3. Mutagenesis on gapped-duplex DNA

A method based on the gapped-duplex approach (Kramer et al., 1984, Nucl. Acids Res. 12, 9441) and a phasmid (plasmid/phage hybrid) was used. Essentially the method rests on a gapped duplex DNA intermediate consisting of a gapped strand (-strand) containing a wild-type antibiotic resistance marker and a template strand (+ strand) carrying an amber mutation in the gene conferring resistance to the antibiotic. After annealing, the mutagenic oligonucleotide becomes incorporated in the gapped strand during in vitro gap-filling and sealing reaction. The resultant molecules are used to transform a mismatch repair deficient (Mut S) host in which the linkage between the intended mutation and the antibiotic resistance marker is preserved. The mixed phasmid population, isolated from this strain, is then allowed to segregate in a suppressor negative host strain. Transformants are plated on antibiotic containing medium, thus imposing a selection for progeny derived from the gapped strand.

The twin vector system pMa/c5-8, which was described by P. Stanssens et al. (Nucl. Acids Res., 1989, 17, 4441) is composed of the following elements:

- pos 11-105 : bacteriophage fd, terminator
- pos 121-215 : bacteriophage fd, terminator
- pos 221-307 : plasmid pBR322 (pos 2069-2153)
- pos 313-768 : bacteriophage f1, origin of replication (pos 5482-5943)
- pos 772-2571 : plasmid pBR322, origin of replication and β -lactamase gene
- pos 2572-2685: transposon Tn903
- pos 2519-2772: tryptophan terminator (double)
- pos 2773-3729: transposon Tn9, chloramphenicol acetyl transferase gene
- pos 3730-3803: multiple cloning site

The sequence is depicted in Figure 1.

In the pMa type vector nucleotide 3409 is changed from G to A, while in the pMc type vector nucleotide 2238 is changed from G to C, creating amber stopcodons in the acetyl transferase gene and β -lactamase gene, respectively, rendering said genes inactive.

All sequences referred to were obtained from Genbank (TM) (release 54), National Nucleic Acid Sequence Data Bank, NIH USA. Plasmid pMc5-8 has been deposited under DSM 4566. To perform mutagenesis the target DNA fragment is cloned into the multiple cloning site of pMa5-8. Subsequently a gapped duplex between pMa5-8 containing the target DNA and pMc5-8 is constructed.

The single strand gap, consisting of the target DNA, can be subjected to mutagenesis with a mutagenic oligonucleotide, with long synthetic oligonucleotides, with a low level of misincorporated nucleotides, with chemicals or with enzymatic misincorporation of nucleotides also random mutagenesis PCR can be applied. For a detailed description see Ausubel et al., *ibid.* or Perbal, *ibid.* As an alternative to in vitro mutagenesis one can use in vivo mutagenesis either with the aid of UV-light or chemicals or by the application of an *E. coli* mutator strain (Fowler et al., J. Bacteriol., 1986, 167, 130).

Mutagenic nucleotides can be synthesised using apparatus obtainable from Applied Bio Systems.

55

4. Random mutagenesis by enzymatic misincorporation of nucleotides

A pMa/pMc gapped duplex can be subjected to primer extension and misincorporation mutagenesis as

originally described by Shortle et al. (Proc. Natl. Acad. Sci. USA, 1982, 79, 1588) by B.C. Cunningham and J.A. Wells (Prot. Eng., 1987, 1, 319) a modification of this procedure is described by Lehtovaara et al. (prot. Eng., 1988, 2, 63).

This method is based on controlled use of polymerases. Four populations of DNA molecules are first generated by primer elongation of a gapped duplex of pMa/pMc so that they terminate randomly, in the gap, but always just before a known type of base (before A, C, G or T, respectively). Each of four populations is then mutagenized in a separate misincorporation reaction where the correct base can now be omitted. In this way all types of base substitution mutations can be generated at every position of the gap. The use of sequenase (TM) (U.S. Biochemical Corporation) was preferred to the use of Klenow polymerase. Moreover MoMuLV reverse transcriptase was used instead of A.M.V. reverse transcriptase, which was used by Lehtovaara et al. (ibid).

To ensure single site substitutions we have introduced the following modification to the protocol described by Lehtovaara et al., ibid. In the reverse transcriptase buffer not three but only one misincorporating nucleotide is present. For instance the A-specific limited base elongation mixture is incubated in three separate reactions with 250 μ M dCTP, 250 μ M dGTP and 250 μ M dTTP, respectively. For a complete set of 4 base specific limited elongation mixtures a total set of 12 separate misincorporation reactions is carried out. After 1.5 hour incubation at 42°C a chase of all four deoxynucleotides in a concentration of 0.5 mM is added and the reactions are further incubated for at least 20 minutes at 37°C. Samples are then further processed according to Lehtovaara et al. (ibid.), with the modification that no counterselection to an uracil-containing DNA strand but a counterselection based on the pMa/c vector was applied.

5. Production of mutant α -amylases

Transformants of E. coli strain WK6 (Zell, R. and Fritz, H.J., EMBO J., 1987, 6, 1809), containing an expression vector, harboring any one of the α -amylase constructs, were inoculated in TB medium (10 ml) at 30°C. TB medium consisted of 0.017M KH_2PO_4 , 0.072M K_2HPO_4 , 12 g/l Bactotryptone, 24 g/l Bacto yeast extract, 0.4% glycerol and an antibiotic (ampicillin with pMa or chloramphenicol with pMc constructs). Samples of the culture were used to inoculate 250 ml TB in 2 liter flasks. At an OD_{600} of 10 - 12, 0.1 mM IPTG (isopropyl- β -D-thiogalactopyranoside) was added and incubation continued for another 12 - 16 hours.

6. Purification of mutant α -amylases

The cells were harvested by centrifugation and resuspended in buffer containing 20% sucrose at 0°C. After a second centrifugation the cells were resuspended in cold water. Cell debris was removed by a third centrifugation and the supernatant was brought to pH 8.0 with 20mM TRIS buffer. CaCl_2 was added to a final concentration of 50mM. The material was heat-treated for 15 min. at 70°C and the insoluble material removed by centrifugation. The supernatant was filtered through 0.22 μ Millipore filter and concentrated to 1/10th of the starting volume.

Further purification was achieved using gelfiltration (on TSK HW-55- Merck) and Mono Q chromatography. Before chromatography on Mono S the pH, of the enzymatic activity containing fractions, was adjusted to 4.8 using sodium acetate. α -amylase was eluted with 250mM NaCl. To avoid inactivation the pH was immediately adjusted to 8.0.

Examples

Example 1

Molecular cloning of Bacillus licheniformis α -amylase gene

Chromosomal DNA isolated from *Bacillus licheniformis* T5 (EP-A-134048; CBS 470.83) was digested with restriction enzyme Eco RI and ligated into the Eco RI site of pUB110 (Gryczan, T.J., et al., J. Bacteriol., 1978, 134, p 318). The ligation mixture was transformed into *Bacillus subtilis* 1A40 (*Bacillus* Genetic Stock Center). Neomycine resistant colonies were tested for α -amylase production on HI agar plates (DIFCO) supplemented with 0.4 g/l starch (Zulkowsky starch, Merck). After growth and incubation with I₂ vapor, a positive colony producing a large clearing halo was selected for further characterization. The plasmid isolated from this positive colony was shown to contain a 3.4 kb Eco RI-Eco RI fragment originating from *Bacillus licheniformis* T5. This plasmid was named pGB33 (EP-A-134048; CBS 466.83). The α -amylase encoding insert was ligated to a synthetic Shine-Dalgarno sequence and the bacteriophage SPO2 promoter resulting in plasmid pProm SPO₂ (see EP-A-0224294; CBS 696.85). The nucleotide sequence of the insert of pProm SPO₂ as determined by the method of Sanger (Proc. Natl. Acad. Sci. U.S.A., 1977, 74, 6463) is shown in Figure 2. The sequence shows a single large open reading frame encoding an α -amylase, which is virtually identical to the α -amylase sequence of *Bacillus licheniformis* as determined by Yuuki et al. (ibid). The first 29 amino acids are a signal sequence which is cleaved off during secretion of the α -amylase. Numbering of amino acids throughout this application refers to the numbering according to the mature protein.

The Yuuki sequence differs at the following positions: at position 134 an Arg is present instead of Leu; at position 310 a Ser is present instead of Gly; at position 320 an Ala is present instead of Ser.

Example 2

Construction of mutagenesis/expression vectors pMaTLia6

Plasmid pPROM SPO₂ was digested with Eco RI and Bcl I and the 1.8 kb Eco RI-Bcl I insert was purified and cloned into Eco RI-Bam HI digested pMa5-8. This pMa5-8 vector was beforehand provided with a modified multiple cloning site. The Bam HI-Hin dIII fragment running from position 3767 to position 3786 in Figure 1 was exchanged for a synthetic DNA sequence as it reads from position 5647 to 5660 in Figure 3. This was carried out to render some restriction sites within the α -amylase gene unique. The resulting α -amylase containing pMa5-8 derivative was digested with Eco RI and Bam HI and ligated to a synthetic DNA fragment carrying a copy of the TAC promoter (De Boer et al., Proc. Natl. Acad. Sci. U.S.A., 1983, 80, 21). The sequence of this synthetic DNA fragment is depicted together with the final α -amylase mutagenesis/expression vector pMaTLia6 in Figure 3 from position 3757 to position 3859. This final α -amylase mutagenesis/expression vector was completed by the introduction of several silent restriction sites which are intended to produce gaps in the α -amylase gene during mutagenesis experiments (Figure 4). For this purpose the following mutations have been made using site-directed oligonucleotide mutagenesis:

- a SpeI site has been introduced by a silent mutation:

T49T

ACG --> ACT

and

S50S

45 AGC --> AGT

- a NarI site has been introduced by the silent mutation:

A269A

GCG --> GCC

- A BstE II site has been introduced just downstream from the TAG stop codon

50 TAGAAGAGC --> TAGGTGACC

This α -amylase mutagenesis vector pMaTLia6 is suited for mutagenesis with the gapped duplex method. Double stranded pMaTLia6 DNA prepared by digestion of suitable restriction enzymes has been annealed to single stranded pMcTLia6 DNA.

The resulting single stranded gaps have been subjected to site-directed mutagenesis, to chemical mutagenesis and to random enzymatic mutagenesis as described in the experimental section.

The availability of the TAC promoter in front of the α -amylase gene enables the inducible expression of α -amylase in *E. coli* by addition of IPTG.

Plasmid pMaTLia6 in *E. coli* WK6 was deposited as CBS 255.89 on June 2nd, 1989.

Example 3Construction of a Bacillus/E. coli shuttle vector for mutagenesis and expression

This vector enables mutagenesis of an inserted gene in *E. coli* and immediate expression in *Bacillus*. The strategy chosen for the construction of the vector was to combine a pUB110 derivative (Gryczan, *ibid.*) with the pMa/c twin vector system in such a way that:

1. The *B. subtilis* cassette can be removed by a single restriction/religation experiment.
2. Different α -amylase genes and different promoters can be easily cloned in this vector.
3. After recircularisation the cloned gene will be under control of a suitable *Bacillus* promoter.
4. During mutagenesis in *E. coli* the *Bacillus* promoter and the structural α -amylase gene are physically separated preventing a possible lethal accumulation of α -amylase in *E. coli*.

A schematic drawing of the shuttle vector is shown in Figure 5. The structure of the final version of the vector pBMa/c1 is depicted in Figure 6. Vector pBMa1 has been deposited under number CBS 252.89, on June 2nd, 1989. The vector has been constructed as follows:

-The Eco RI-Sna BI fragment of pUB110 carrying the REP-gene and the Neo^R gene was purified and cloned into Eco RI-Sma I digested pUC8.

- The Eco RI-Hin dIII fragment of this pUC8 derivative was cloned into Eco RI-Hin dIII digested pMa5-8 resulting in plasmid pMa5-80.

- The Bam HI-Xba I polylinker fragment was substituted by a synthetic fragment of DNA encoding the SPO₂ promoter of bacteriophage SPO₂ (Williams et al., J. Bacteriol., 1981, 146, 1162) plus restriction recognition sites for Sac II, Apa I, Xho I, Sac I, Bgl I, Mlu I and Xba I.

- The unique Eco RI site of pMa5-80 was used to insert a polylinker fragment constituting the following recognition sites: Eco RI, Sma I, Sac I, Eco RV, Sph I, Kpn I, Xba I and Hin dIII.

For specific purposes derivatives pBMa/c2 and pBMa/c6 have been developed out of pBMa/c1.

- In pBMa/c2 the Eco RI-Hin dIII polylinker of pBMa/c1 has been replaced by the corresponding polylinker of pUC19.

- In pBMa/c6 in addition the Sac II site in the right polylinker of pBMa/c1 has been removed by a Klenow reaction.

Site directed mutagenesis on the *B. licheniformis* α -amylase gene was performed after construction of pBMa/c6 Lia6. This vector was constructed by ligating the Bam HI-Hin dIII fragment isolated from pMaTLia6 into the above mentioned pBMa/c6 which was cleaved by Bam HI and Hin dIII. The resulting plasmid (Figure 7) can be used to construct gapped duplexes for mutagenesis in *E. coli*.

The resulting mutants have been expressed in *Bacillus subtilis* 1A40 (BGSC 1A40) after restriction with Sac I, religation and transformation according to Chang and Cohen (Mol. Gen. Genet., 1979, 168, 111).

Example 4Expression in E. coli of correctly matured Bacillus licheniformis α -amylase

Characterization of the α -amylase produced by pMaTLia 6 (Example 2) showed that a portion of the α -amylase was incorrectly processed during secretion. NH₂-terminal sequencing revealed an extra Alanine residue for α -amylase produced in *E. coli* WK 6.

Although we have no indication that this will give different properties to the amylase we have replaced the α -amylase signal sequence by the alkaline phosphatase Pho A signal sequence. To this end a mutagenesis experiment was carried out so as to introduce a Fsp I restriction site in pMaTLia 6 at the junction of the signal peptide and the mature α -amylase. After Fsp I and Bam HI digestion a synthetic DNA fragment encoding the pho A signal sequence (Michaelis et al., J. Bacteriol., 1983, 154, 366) was inserted. The sequence of this construction is shown in Figure 8. α -Amylase produced by pMaCTPLia6 was shown to possess the correct NH₂-terminal sequence.

A. Screening for acid-stable α -amylase mutants

Method:

Possible mutants are grown in microtiterplates. The growth medium is 250 μ l Brain Heart Infusion broth (DIFCO). The following additions are made:

chloramphenicol	50 μ g/ml
I.P.T.G. (SIGMA)	0.2 mM
CaCl ₂	2 mM

These microtiterplates are placed at 37° C for 40 hours without shaking.

After this time period, in which the α -amylase is produced, 5 μ l samples are taken from each well and spotted on 2 different types of agar plates (144 x 140 mm). The first type is a rich Heart-Infusion agar plate (DIFCO) + 0.4% starch (Zulkowsky starch-Merck) + chloramphenicol 50 μ g/ml. After incubation at 37° C for 16 hours this plate serves as a storage for mutants.

The second type of plate is the actual screening plate, it contains:

Bacto agar (DIFCO)	1.5%
Zuikowsky starch	0.2%

Agar and starch are dissolved in synthetic tap water (STW). This is: demineralised water +

CaCl ₂	2 mM
MgCl ₂	1 mM
NaHCO ₃	2.5 mM
BSA	10 µg/ml

11

pH values in the agar plate when measured are somewhat lower than those of the stock solutions. From each well 5 μ l of culture is spotted on 3 screening plates with different pH values.

The pH-range is chosen in such a way that there is little or no activity left for the wild-type α -amylase on the plate with the lowest pH-value.

3. Colouring

The screening plates are incubated for 2 hours at 55° C. After this period an I₂ solution is poured over the plates. 10 x I₂ solution contains 30 g I₂ and 70 g KI per liter.

The amount of clearance of the spots is correlated with the residual α -amylase activity at that pH value. Those mutants that perform better than the wild-type controls are selected for a second round of screening. Wild-type halo's are very reproducible in this experiment.

4. Second screening

Positive mutants are picked from the rich plate and purified on fresh HI plates + chloramphenicol. 4 single colonies are picked from each mutant and they are tested again in a similar way as in the first screening. In addition serial dilutions of these cultures are made with STW and these dilutions are spotted on neutral pH screening plates (pH = 7.0). Comparison with wild-type cultures enables one to decide if the better performance at low pH is due to an overall better α -amylase production or to intrinsically more stable α -amylase.

The mutants that "survive" the second screening are characterized by determining the nucleotide sequence of that part of the gene that was subjected to mutagenesis.

B. Screening for alkali stable α -amylase

Screening for alkali stable α -amylases is performed in a manner similar to the one used for acid stable α -amylase. After growth in microtiter plates 5 μ l samples are taken from each well and spotted onto a storage plate and onto the actual screening plate. The latter is composed of:

Bacto Agar (DIFCO)	1.5%
Zulkowsky starch	0.2%

and completed with demineralized water plus

CaCl ₂	2 mM
MgCl ₂	1 mM
NaHCO ₃	2.5 mM
BSA	10 μ g/ml

The screening plates are buffered with 50 mM carbonate/bicarbonate buffer, pH values are 9.0, 9.5 and 10.0. The pH range is chosen in such a way that there is little or no activity of the wild-type α -amylase at the highest pH value. After 2 hours incubation at 55° C an I₂ solution is poured over the plates. Those mutants that give a better halo than the wild-type enzyme are selected for a second round of screening. This second round of screening is performed in a similar fashion as the screening for the acid stability.

C. Screening for thermostable α -amylase mutants

α -Amylase mutants that perform better or worse at high temperature than the wild-type α -amylase, can also be selected by comparison of halo's on starch plates caused by the residual amylase activity in the

culture broths after heating.

Method:

5

1. Mutants are grown in the same way as for the pH screening.
2. The mutants are replicated on HI agar plates as for the pH screening.
3. The separate wells of the microtiterplates were closed with disposable caps (Flow laboratories) to prevent evaporation of the culture broths during the heating step.
4. Microtiterplates were heated in a waterbath for 1 hour at 95 °C. After heating the microtiterplates were placed in a centrifuge for collecting the total sample on the bottom of the microtiterplate.
5. Screening for thermostable mutants was done as follows:

10

From each well 5 µl of culture was spotted on neutral screeningplates (See pH-screening). These plates were incubated for 1 hour at 55 °C.

15

After staining the starch with the iodine solution mutants and controls can be screened for residual α -amylase activity by comparing clearance of the spots (halo's).

In case the residual activity of the controls is too high, serial dilutions must be made and spotted on the screening plate to be able to discriminate for mutants that are more thermostable than the wild-type enzyme.

20

6. Possible interesting mutants are tested further as was done in the pH-screening method.

A combination of screening type A or B with type C can be applied if a combination of properties is desired. For instance after the first round of screening for alkali stable α -amylase, a second round of screening for thermostability can be performed. Those mutants that score positive in both tests may be selected as candidates exhibiting a combination of desired properties.

25

Example 6

30

Bisulphite mutagenesis of pMaTLia6

Single stranded DNA of pMaTLia6 was annealed with Sac II-Cla I digested pMcTLia6 in order to obtain a heteroduplex with a gap running from position 4315 to 4569 (Figure 3). This heteroduplex was subjected to bisulphite mutagenesis (see experimental).

35

After transformation into *E. coli* WK6 mut S (Zell, R. and Fritz H.J., *ibid*) and selection on chloramphenicol containing agar plates (50 µg/ml) plasmid pools were isolated and transformed into *E. coli* WK6. *E. coli* WK6 Mut S was deposited as CBS 472.88, *E. coli* WK6 was deposited as CBS 473.88.

40

Resulting transformants were grown in BHI medium (DIFCO) containing 2.0 mM CaCl₂, 50 µg/ml chloramphenicol and 0.20 mM IPTG (SIGMA) during 40 hours at 37 °C in microtiter wells without shaking. Screening for pH stable mutants was carried out as described in Example 5.

45

About 300 Cm^R transformants were screened. The mutation frequency as determined by DNA sequencing was on average 0.4 mutation/molecule over the gap. One acid stable mutant, D7, was identified after the pH screening. Sequencing of this mutant revealed mutation H133Y originating from a mutation of the encoding triplet from CAC to TAC.

Mutant D7 was also found positive in the thermostability screening assay (Example 5).

50

DNA sequencing was performed on single stranded DNA with a specific oligonucleotide designed to prime just before the Sac II-Cla I fragment. In a separate mutagenesis experiment 1000 Cm^R transformants were screened. Another acid stable mutant, 2D5, was identified after the pH screening. This mutant has the following mutations:

H133Y CAC --> TAC

T149I ACA --> ATA

55

Bisulphite mutagenesis has been applied in a similar manner as just described on the Cla I-Sal I gap which runs from position 4569 to position 4976 of Figure 3. About 300 Cm^R transformants were screened (mutation frequency 0.6 mutations/molecule). No acid stable transformants were found. A number of acid labile mutants were found. Among these acid labile mutants some may have a shifted pH spectrum resulting in a more alkaline stable phenotype.

Example 7Enzymatic mutagenesis of pMatLia6

Single stranded pMatLia6 (Figure 4) was annealed with Cla I-Sal I digested pMcTLia6 in order to obtain a heteroduplex running from position 4569 to 4976 (Figure 3). The gapped duplex was subjected to enzymatic misincorporation mutagenesis as described in the experimental section.

A sample obtained after dATP-limited primer elongation was split in three parts and incubated in the presence of reverse transcriptase with dCTP, dGTP and dTTP, respectively. After incubation at 37 °C for 10 minutes a chase with all four dNTP's and Klenow polymerase was given T4-DNA ligase was added to finish the elongation to completely double stranded molecules.

These molecules were transformed into E. coli WK 6 Mut S and plasmid pools were recovered. These plasmid pools were subsequently transformed into E. coli WK 6 and the colonies were selected on chloramphenicol (50 µg/ml) containing agar plates. Resulting mutants were screened for stability of α-amylase as described in Example 5.

In another experiment the Spe I-Sac II gap was subjected to limited primer elongation with dATP, dCTP, dGTP and dTTP, respectively. These primer pools were mutagenized by misincorporation (see experimental). 100 Cm^R transformants were tested on pH plates (Example 5) and mutant M29 was identified as more stable at low pH. The sequence of the mutation was determined: A111T GCG → TCG

Example 8Properties of stable mutants

Two of the mutants obtained from the bisulphite mutagenesis experiments were further characterized. As described before DNA sequencing suggested the following amino acid replacements;

- D7 contained a tyrosine at position 133 instead of a histidine (D7 = H133Y),
- 2D5 contained the D7 mutation and in addition threonine 149 was replaced by isoleucine (2D5 = H133Y, T149I).

a) Measurement of enzymatic activity

The enzymatic activity of B. licheniformis α-amylase WT and mutants was measured using 4-nitrophenyl-maltopentaoside (4NP-DP5) as a substrate, 4 nitrophenol and maltopentaose are formed, this reaction can be followed by measuring the change in OD 405. The assay was performed at 35 °C in 50mM MOPS, 50mM NaCl, 2mM CaCl₂ (PH 7.15) and 0-1mm 4NP-DP5.

Initial rates were measured and E-nitrophenol was taken as 10,000 1/M/cm. Figure 9 shows the results for WT and 2D5 α-amylases. Vmax and Km were calculated and are given in Table 1.

Table 1

	Vmax(µmol/min/mg)	Km(mM)
WT	66.7 ± 0.9	0.112 ± 0.005
2D5	66.3 ± 0.7	0.119 ± 0.004

Table 1 clearly shows that the mutations of α-amylase 2D5 do not influence the enzymatic activity in a substantial way.

b) Influence of Ca^{2+} on the thermoinactivation

Heat inactivation experiments were performed for WT, D7 and 2D5 at varying calcium concentrations. The procedure was as follows :

1) Demetallization

Enzyme (2 - 3 mg/ml) dialyzed for 24 hrs against 3 x 1 L

20 mM MOPS

5 mM EDTA

5 mM EGTA pH 7.0

3 x 1 L

20mM MOPS pH 7.0

2) Remetallization

- 500 μl buffer 100 mM (e.g. MES, MOPS, EPPS)*

- 145 μl demetallized enzyme (e.g. 2.15 mg/ml)

- 100 μl CaCl_2 (100, 50, 30, 20, 10, 5 or 2.5 mM)

- x μl K_2SO_4 (100 mM)

- (255-x) μl H_2O

[CaCl_2] final (mM)	[K_2SO_4] final (mM)
0,25	14,75
0,5	14,5
1	14
2	13
3	12
5	10
10	0

* - pH MES e.g. 6.77 at room temperature will give 6.0 at 90 °C (pKa 6.15 pKa/°C = -0.011)

- pKa were from Table of Merck (Zwitterionische Puffersubstanzen)

3) Heat-inactivation

1 ml enzyme solution preincubated at room temperature was heated at 90.5 °C or 95 °C in closed Pierce-vials (teflon coated-seals) at a concentration of about 0.2 mg/ml. 50 μl samples were withdrawn at regular intervals between 0 and 6 hrs with a syringe and cooled on ice. Residual activities have been determined with 4NP-DP5 (0.5mM).

Half lives were determined using a single exponential decay fitting program (GRAPHPAD).

Figures 10 and 11 show the half life times of WT and D7 α -amylases at pH 5.5 and 7.0 respectively as a function of the Ca^{2+} concentration at 90.5 °C. The Ca^{2+} dependence of 2D5 has only been determined at pH 7.0 at 95 °C (Figure 12). It can also be seen that the Ca^{2+} dependence of the mutants is not different from that of the WT.

c. Thermostability of mutant α -amylases at different pH values

The pH dependence of thermoinactivation for both D7 and 2D5 has been determined at 90.5 and 95 °C respectively using the buffer as described above at a 1 mM Ca^{2+} concentration. It can be concluded that the thermal stability of both D7 and 2D5 is greatly increased (up to twofold for 2D5) over the entire pH range. (Figures 13 and 14).

Example 9Production of mutant enzymes in Bacillus

Mutations in the *B. licheniformis* α -amylase, which were identified by expression in *E. coli* WK6 were transferred to a *Bacillus* expression vector in two different ways.

- 10 a) With the aid of the unique restriction sites within the α -amylase gene (Figure 4), fragments carrying mutations were isolated from pMaTLia6 mutants and subcloned into the homologous position of pBMa6.Lia6. The latter plasmid, which can be replicated either in *E. coli* or in *Bacillus*, was subsequently digested with *Sac* I and recircularized with T4 DNA ligase. After transformation into *Bacillus subtilis* 1A40 high level α -amylase production under control of the *SPO*₂ promoter was obtained.
- 15 Recircularized pBMa6.Lia6 is named pB6.Lia6 to indicate the removal of the *E. coli* portion of the vector.
- b) pBMa6.Lia6 single stranded DNA was recollected from *E. coli* and annealed with restriction enzyme digested pBMc6.Lia6 double stranded DNA in order to obtain a gapped duplex with the intended gap on the α -amylase gene. This gap was then subjected to site-directed mutagenesis with an oligonucleotide (as described in the experimental section) which encodes the desired mutation. pBMc6.Lia6 vector is
- 20 then transformed into pB6.Lia6 type vector as described above. Combination of different single site mutation can be performed by method a) if mutations are in different gaps, preferably, however, method b) is used.

The mutations of mutants D7 and 2D5 were transferred to pBMa6.Lia6 by method a) by exchanging the *Sac* II-*Sal* I fragments and α -amylase was recovered from the medium of transformed *Bacillus subtilis* 1A40.

25 Supernatants of both mutants were subjected to the screening procedures of Examples and it was confirmed that both mutants produce α -amylase which is more acid stable and more thermostable than α -amylase produced by wild-type pB6.Lia6.

The phenotype of the α -amylase mutations in *Bacillus* is thus not different from the phenotype in *E. coli*.

30 Ultimately pB6.Lia6 mutants have been transformed into *Bacillus licheniformis* T9, which is a protease negative, α -amylase negative derivative of *Bacillus licheniformis* T5, (EP-0253455, CBS 470.83). Host T9 has been used to produce high level amounts of α -amylase mutants in a homologous system. The removal of the chromosomal α -amylase gene renders this strain very suited for the production of mutant α -amylase as no contaminating wild-type α -amylase is being produced anymore. Enzyme recovered from this strain

35 has been used for industrial application testing. The industrial use of mutants pB6.Lia6.2D5 and pB6.Lia6.D7 was demonstrated.

Example 10Application test of mutant α -amylase under conditions of starch liquefaction

45 To test mutant α -amylase 2D5 in more realistic circumstances, we have purified the fermentation broth (of Example 9) with ultrafiltration and formulated the enzyme with 50% propylene glycol. Three samples have been tested:

50

893701 : WT	<i>B. licheniformis</i> T5 α -amylase	1530 TAU/g
893703 : 2D5	Mutant prepared as WT	2820 TAU/g
Maxamyl 0819	Commercial sample	7090 TAU/g

55 One TAU (thermostable α -amylase unit) is defined as the quantity of enzyme that will convert under standardized conditions 1 mg of starch per minute in a product having an equal absorption to a reference colour at 620 nm after reaction with iodine. Standard conditions are pH 6.6; 30 °C; reaction time : 20 min.

Reference colour is 25g CoCl₂ · 6H₂O, 3.84 g K₂Cr₂O₇ and 1 ml HCl (1M) in 100 ml distilled H₂O.

1. Liquefaction test at low pH (5.5 and 5.25)

The temperature of starch slurry is increased to $110 \pm 0.5^\circ \text{C}$ as quick as possible and kept at this temperature for 6 minutes.

The liquefaction is realized in continuous flow (5.4 l/h). 3 Samples of 135 ml (1.5 minute of liquefaction) are taken after 45, 60 and 75 minutes of liquefaction and kept at 95°C for two hours. After this time, 50 ml of the sample are acidified with 0.4 ml H₂SO₄ N to obtain pH 3.5 and put in boiling bath for 10 minutes in order to stop enzymatic activity before D.E. determination.

The remaining part of the sample is cooled in order to determine residual enzymatic activity.

Slurry composition:

3.3 kg corn starch D.S. 88% (2.904 kg dry starch).

5.45 l well water (40 T.H.).

Dry substance of the slurry is 33%.

pH is corrected at 5.5 with 1N sulfuric acid or 1N NaOH.

Enzyme concentration: 4.4 TAU/gr dry starch.

The flow rate is verified two or three times during the trial.

2. Determination of D.E.

Dry substance of liquefied starch is verified with a refractometer (about 34%). D.E. is determined with the well-known Lane Eynon method. The results are shown in Figure 15.

3. Residual Enzymatic Activity

Residual amylase activity in liquefied starch is determined with a Brabender amylograph.

40 g potato starch

390 ml distilled water at 50°C

50 ml Tris buffer 0.05 M pH 6.50

5 ml CaCl₂ 2H₂O at 30 g/l

The temperature is increased to 80°C ($1.5^\circ/\text{min}$) when viscosity is stabilized (10 min) 5 ml of diluted liquefied starch (7 g up to 50 ml with distilled water) is added, the decrease of viscosity after 20 minutes is measured, this decrease is a function of the enzymatic activity. A standard curve with known enzymatic concentration allows to estimate residual activity in T.A.U.

Mutant 2D5 performs significantly better at pH < 5.5 and 110°C than WT enzyme. An improvement of 2-3 DE units at pH 5.25 is obtained with mutant 2D5.

Example 11

Application test of mutant α -amylase under conditions of textile desizing

To test the industrial application of alkaline α -amylase mutants a test is performed on the stability at 20°C in the following solution:

1.4% H₂O₂ (35%)

1.0-1.5% Caustic Soda (100%)

15-20 ml/l Sodium Silicate (38 Bé)

0.3-0.5% Alkylbenzene sulphonate (Lanaryl N.A.-ICI)

0.5-1.0% Organic stabilizer (Tinoclarite G)

After incubation during 2.5 hours the α -amylase mutants selected for their desired properties should have any remaining enzyme activity.

Claims

1. A mutant α -amylase, that is the expression product of a mutated DNA sequence encoding an α -amylase, characterized in that the mutant α -amylase has an amino acid sequence which differs at least in one amino acid from the wild-type enzyme and that said mutant α -amylase exhibits improved properties for application in the degradation of starch and/or textile desizing wherein the improved properties are due to the amino acid replacements.
2. An α -amylase according to Claim 1, characterized in that it exhibits improved thermostability.
3. An α -amylase according to Claim 1, characterized in that it exhibits improved stability at a pH below 6.5 and/or above 7.5.
4. An α -amylase according to Claim 1, characterized in that it exhibits improved thermostability and acid stability.
5. An α -amylase according to any one of the Claims 1-4, in which the original gene from which the mutant enzyme is derived is obtained from a microorganism, preferably a Bacillus strain.
6. An α -amylase according to Claim 5, in which said gene is derived from a wild-type gene of a strain selected from the group consisting of B. stearothermophilus, B. licheniformis and B. amyloliquefaciens.
7. An α -amylase according to Claim 6, characterized in that this enzyme differs from the wild-type α -amylase obtainable from Bacillus licheniformis by an amino acid replacement at one or more of the positions 111, 133 and 149 or at corresponding positions in any homologous α -amylase.
8. An α -amylase according to Claim 7, characterized in that it contains one or more of the following amino acid replacements: Ala-111-Thr, His-133-Tyr, Thr-149-Ile.
9. A mutant gene encoding an α -amylase as defined in any one of Claims 1-8.
10. An expression vector which comprises a mutant gene according to Claim 9.
11. A host cell harboring an expression vector according to Claim 10.
12. A host cell which is substantially incapable of producing extracellular amylolytic enzymes prior to transformation, characterized in that it is transformed with an expression vector according to Claim 10.
13. A host cell according to Claim 12 being B. licheniformis T9.
14. A Bacillus / E. coli shuttle vector, wherein the expression of the cloned gene in E. coli is made impossible by physical separation of the regulatory sequences from the structural gene and wherein the expression of the cloned gene in Bacillus can be restored by digestion with a single restriction enzyme and subsequent recircularization.
15. A method for preparing an amylolytic enzyme having improved properties for application in starch degradation or in textile desizing which comprises the following steps:
mutagenizing a cloned gene encoding an amylolytic enzyme of interest or a fragment thereof;
isolating the obtained mutant amylase gene or genes;
introducing said mutant amylase gene or genes into a suitable host strain for expression and production;
recovering the produced mutant amylase and identifying those mutant amylases having improved properties for application in starch degradation or textile desizing.
16. A process for producing a mutant α -amylase comprising;
- cultivating a host cell according to any of Claims 11-13 in a suitable medium,
- recovering the produced α -amylase.
17. Use of the α -amylase according to any one of the Claims 1-8 in starch degradation and in textile desizing.
18. Process for the degradation of starch which comprises the use of a mutated α -amylase according to any one of the Claims 1-8.
19. Process for textile desizing which comprises the use of a mutated α -amylase according to any one of the Claims 1-8.
20. Starch degradation composition comprising a mutated α -amylase according to any one of the Claims 1-8.
21. Textile desizing composition comprising a mutated α -amylase according to any one of the Claims 1-8.

```

      10      20      30      40      50      60
AATTCACCTCGAAAGCAAGCTGATAAACCGATACAATTAAGGCTCCTTTTGGAGCCTTT

      70      80      90     100     110     120
TTTTTTGGAGATTTTCAACGTGAAAAAATTATTATTCGCAATTCCAAGCTAATTCACCTC

      130     140     150     160     170     180
GAAAGCAAGCTGATAAACCGATACAATTAAGGCTCCTTTTGGAGCCTTTTTTTTGGAG

      190     200     210     220     230     240
ATTTTCAACGTGAAAAAATTATTATTCGCAATTCCAAGCTCTGCCTCGCGCGTTTCGGTG

      250     260     270     280     290     300
ATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTGTAAG

      310     320     330     340     350     360
CGGATGCAGATCACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGGGTGTGGTGGTTACGC

      370     380     390     400     410     420
GCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTCGCTTCTTCCCTT

      430     440     450     460     470     480
CCTTCTCGCCACGTTTCGCCGGCTTTCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAG

      490     500     510     520     530     540
GGTTCGGATTTAGTGCTTTACGGCACCTCGACCCCAAAAACTTGATTAGGGTGATGGTT

      550     560     570     580     590     600
CACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGT

      610     620     630     640     650     660
TCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAACCCTATCTCGGTCTATT

      670     680     690     700     710     720
CTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTT

      730     740     750     760     770     780
AACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTACAATTTGATCTGCGCTCG

      790     800     810     820     830     840
GTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACA

      850     860     870     880     890     900
GAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAAC

      910     920     930     940     950     960
CGTAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC

      970     980     990    1000    1010    1020
AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCG

      1030    1040    1050    1060    1070    1080
TTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATAC

      1090    1100    1110    1120    1130    1140
CTGTCCGCCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCAATGCTCACGCTGTAGGTAT

```

Figure 1

1150	1160	1170	1180	1190	1200
CTCAGTTCGGTGTAGGTGCTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAG					
1210	1220	1230	1240	1250	1260
CCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGAC					
1270	1280	1290	1300	1310	1320
TTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGCCGGT					
1330	1340	1350	1360	1370	1380
GCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGT					
1390	1400	1410	1420	1430	1440
ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGC					
1450	1460	1470	1480	1490	1500
AAACAAACCCGCTGGTAGCCGGTGGTTTTTTTGGTTTGCAAGCAGCAGATTACGCGCAGA					
1510	1520	1530	1540	1550	1560
AAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAAC					
1570	1580	1590	1600	1610	1620
GAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATC					
1630	1640	1650	1660	1670	1680
CTTTTAAATTAAAAATGAAGTTTTTAAATCAATCTAAAGTATATATGAGTAAACTTGGTCT					
1690	1700	1710	1720	1730	1740
GACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTCA					
1750	1760	1770	1780	1790	1800
TCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCT					
1810	1820	1830	1840	1850	1860
GGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCA					
1870	1880	1890	1900	1910	1920
ATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCTCC					
1930	1940	1950	1960	1970	1980
ATCCAGTCTATTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCGCCAGTTAATAGTTTG					
1990	2000	2010	2020	2030	2040
CGCAACGTTGTTGCCATTGCTGCAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCT					
2050	2060	2070	2080	2090	2100
TCATTACAGCTCCGGTCCCAACGATCAAGGCGAGTTACATGATCCCCCATGTTGTGCAAA					
2110	2120	2130	2140	2150	2160
AAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAAGTAAGTTGGCCGCAAGTGA					
2170	2180	2190	2200	2210	2220
TCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATGC					
2230	2240	2250	2260	2270	2280

Fig. 1 (continued)

TTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGGCACC
 2290 2300 2310 2320 2330 2340
 AGTTGCTCTTGGCCGGCGTCAACACGGGATAATACCGCGCCACATAGCAGAACTTTAAAA
 2350 2360 2370 2380 2390 2400
 GTGCTCATCATTGAAAAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGCTGTTG
 2410 2420 2430 2440 2450 2460
 AGATCCAGTTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTTACTTTC
 2470 2480 2490 2500 2510 2520
 ACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGG
 2530 2540 2550 2560 2570 2580
 GCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCAGACAG
 2590 2600 2610 2620 2630 2640
 TTTTATTGTTTCATGATGATATATTTTATCTTGTGCAATGTAACATCAGAGATTTTGAGA
 2650 2660 2670 2680 2690 2700
 CACAACGTGGCTTTGTTGAATAAATCGAACTTTTGTGAGTTGACTCCCCGCGCGCGATG
 2710 2720 2730 2740 2750 2760
 GGTCCAATTTGCTTTCGAAAAAAAAGCCCCGCTCATTAGGCGGGCTAAAAAAAAGCCCCGCT
 2770 2780 2790 2800 2810 2820
 CATTAGGCGGGCTCGAATTTCTGCCATTATCCGCTTATTATCACTTATTCAGGCGTAGC
 2830 2840 2850 2860 2870 2880
 AACCAGGCGTTTAAGGGCACCAATAACTGCCTTAAAAAATTACGCCCGCCCTGCCACT
 2890 2900 2910 2920 2930 2940
 CATCGCAGTACTGTTGTAATTCATTAAGCATTCTGCCGACATGGAAGCCATCACAGACGG
 2950 2960 2970 2980 2990 3000
 CATGATGAACCTGAATCGCCAGCGGCATCAGCACCTTGTCGCCTTGCGTATAATATTGCG
 3010 3020 3030 3040 3050 3060
 CCATAGTGAAAACGGGGGCGAAGAAGTTGTCCATATTCGCCACGTTTAAATCAAACTGG
 3070 3080 3090 3100 3110 3120
 TGAAACTCACCCAGGGATTGGCTGAGACGAAAAACATATTCTCAATAAACCTTTAGGGA
 3130 3140 3150 3160 3170 3180
 AATAGGCCAGGTTTTACCGTAACACGCCACATCTTGCGAATATATGTGTAGAACTGCC
 3190 3200 3210 3220 3230 3240
 GGAAATCGTCTGGTATTCACCTCCAGAGCGATGAAAACGTTTCAGTTTGCTCATGGAAAA
 3250 3260 3270 3280 3290 3300
 CGGTGTAACAAGGGTGAACACTATCCCATATCACCAGCTCACCGTCTTTCATTGCCATAC
 3310 3320 3330 3340 3350 3360
 GAAATTCGGGATGAGCATTTCATCAGGCGGGCAAGAATGTGAATAAAGGCCGGATAAACT

Fig. 1 (continued)

3370 3380 3390 3400 3410 3420
TGTGCTTATTTTTCTTTACGGTCTTTAAAAAGGCCGTAATATCCAGCTAAACGGTCTGGT

3430 3440 3450 3460 3470 3480
TATAGGTACATTGAGCAACTGACTGAAATGCCTCAAAATGTTCTTTACGATGCCATTGGG

3490 3500 3510 3520 3530 3540
ATATATCAACGGTGGTATATCCAGTGATTTTTTTCTCCATTTTAGCTTCCTTAGCTCCTG

3550 3560 3570 3580 3590 3600
AAAATCTCGATAACTCAAAAAATACGCCCGGTAGTGATCTTATTTTATTATGGTGAAAGT

3610 3620 3630 3640 3650 3660
TGGAACCTCTTACGTGCCGATCAACGTCTCATTTTTCGCCAAAAGTTGGCCCAGGGCTTCC

3670 3680 3690 3700 3710 3720
CGGTATCAACAGGGACACCAGGATTTATTTATTCTGCGAAGTGATCTTCCGTCACAGGTA

3730 3740 3750 3760 3770 3780
TTTATTCTGAAGACGAAAGGGCATCGCGCGCGGGAATCCCGGGGATCCGTCGACCTGCA

3790 3800
GCCAAGCTTGGTCTAGAGGTCTGA

10 20 30 40 50 60
 GTCTACAAACCCCTTAAAAACGTTTTTAAAGGCTTTTAAAGCCGCTGTGTACGTTCTTAAAG
 70 80 90 100 110 120
 GAATTTCACACTGGCCTTGGTTAAGGTTAAGATGTGGACGGAATCGGTAAAGTGTAGTAAA
 130 140 150 160 170 180
 GTACAATTAATCGGGAGCTTAGATGTCCCTTCAACATCTTATATAGAAGGGAAGGTTGGC
 190 200 210 220 230 240
 AAATGGAAATTGAAAGAATTAACGAGCATACAGTAAAAATTTTATATGTCTTACGGAGATA
 250 260 270 280 290 300
 TTGAAGATCGCGGTTTTGACAGAGAAGAAATTTGGTATAACCGTGAGCCGAGTGAAGAAC
 310 320 330 340 350 360
 TTTTCTGGGAAGTCATGGATGAAGTTCATGAAGAAGAGGAATTCGAGCTCGCCCCGGGGAT
 370 380 390 400 410 420
 CCAAGGAGGTGATCTAGAGTCATGAAACAACAAAAACGGCTTTACGCCCCGATTGCTGACG
 M K Q Q K R L Y A R L L T
 430 440 450 460 470 480
 CTGTTATTTGCGCTCATCTTCTTGCTGCCTCATTCTGCAGCAGCGGCGGCAAATCTTAAT
 L L F A L I F L L P H S A A A A A N L N
 +1
 490 500 510 520 530 540
 GGGACGCTGATGCAGTATTTTGAATGGTACATGCCCAATGACGGCCAACATTGGAAGCGT
 G T L M Q Y F E W Y M P N D G Q H W K R
 5
 550 560 570 580 590 600
 TTGCAAAACGACTCGGCATATTTGGCTGAACACGGTATTACTGCCGTCTGGATTCCCCCG
 L Q N D S A Y L A E H G I T A V W I P P
 25
 610 620 630 640 650 660
 GCATATAAGGGAACGAGCCAAGCGGATGTGGGCTACGGTGCTTACGACCTTTATGATTTA
 A Y K G T S Q A D V G Y G A Y D L Y D L
 45
 670 680 690 700 710 720
 GGGGAGTTTCATCAAAAAGGGACGGTTCGGACAAAAGTACGGCACAAAAGGAGAGCTGCAA
 G E F H Q K G T V R T K Y G T K G E L Q
 65
 730 740 750 760 770 780
 TCTGCGATCAAAAGTCTTCAATTCGCGGACATTAACGTTTACGGGGATGTGGTCATCAAC
 S A I K S L H S R D I N V Y G D V V I N
 85
 790 800 810 820 830 840
 CACAAAGGCGGCGCTGATGCGACCGAAGATGTAACCGCGGTTGAAGTCGATCCCGCTGAC
 H K G G A D A T E D V T A V E V D P A D
 105
 850 860 870 880 890 900
 CGCAACCGCGTAATTTTCAGGAGAACACCTAATTAAGCCTGGACACATTTTCATTTTCCG
 R N R V I S G E H L I K A W T H F H F P
 125
 910 920 930 940 950 960
 GGGCGCGGCAGCACATACAGCGATTTTAAATGGCATTGGTACCATTTTGACGGAACCGAT
 G R G S T Y S D F K W H W Y H F D G T D
 145

Figure 2

970 980 990 1000 1010 1020
 TGGGACGAGTCCCGAAAGCTGAACCGCATCTATAAGTTTCAAGGAAAGGCTTGGGATTGG
 W D E S R K L N R I Y K F Q G K A W D W
 165

1030 1040 1050 1060 1070 1080
 GAAGTTTCCAATGAAAACGGCAACTATGATTATTTGATGTATGCCGACATCGATTATGAC
 E V S N E N G N Y D Y L M Y A D I D Y D
 185

1090 1100 1110 1120 1130 1140
 CATCCTGATGTCGCAGCAGAAATTAAGAGATGGGGCACTTGGTATGCCAATGAACTGCAA
 H P D V A A E I K R W G T W Y A N E L Q
 205

1150 1160 1170 1180 1190 1200
 TTGGACGGTTTCCGTCTTGATGCTGTCAAACACATTAAATTTCTTTTTTGC GG GATTGG
 L D G F R L D A V K H I K F S F L R D W
 225

1210 1220 1230 1240 1250 1260
 GTTAATCATGTCAGGGAAAAACGGGGAAGGAAATGTTTACGGTAGCTGAATATTGGCAG
 V N H V R E K T G K E M F T V A E Y W Q
 245

1270 1280 1290 1300 1310 1320
 AATGACTTGGGGCGCGCTGGAAAACTATTGAAACAAAACAAATTTTAATCATTTCAGTGT
 N D L G A L E N Y L N K T N F N H S V F
 265

1330 1340 1350 1360 1370 1380
 GACGTGCCGCTTCATTATCAGTTCATGCTGCATCGACACAGGGAGGCGGCTATGATATG
 D V P L H Y Q F H A A S T Q G G G Y D M
 285

1390 1400 1410 1420 1430 1440
 AGGAAATTGCTGAACGGTACGGTCGTTTCCAAGCATCCGTTGAAATCGGTTACATTTGTC
 R K L L N G T V V S K H P L K S V T F V
 305

1450 1460 1470 1480 1490 1500
 GATAACCATGATACACAGCCGGGGCAATCGCTTGAGTCCGACTGTCCAAACATGGTTTAAG
 D N H D T Q P G Q S L E S T V Q T W F K
 325

1510 1520 1530 1540 1550 1560
 CCGCTTGCTTACGCTTTTATTCTCACAAGGGAATCTGGATACCCTCAGGTTTTCTACGGG
 P L A Y A F I L T R E S G Y P Q V F Y G
 345

1570 1580 1590 1600 1610 1620
 GATATGTACGGGACGAAAGGAGACTCCCAGCGCGAAATTCCTGCCTTGAAACACAAAATT
 D M Y G T K G D S Q R E I P A L K H K I
 365

1630 1640 1650 1660 1670 1680
 GAACCGATCTTAAAAGCGAGAAAACAGTATGCGTACGGAGCACAGCATGATTATTTTCGAC
 E P I L K A R K Q Y A Y G A Q H D Y F D
 385

1690 1700 1710 1720 1730 1740
 CACCATGACATTGTGGGCTGGACAAGGGAAGGCGACAGCTCGGTTGCAAATTCAGGTTTG
 H H D I V G W T R E G D S S V A N S G L
 405

1750 1760 1770 1780 1790 1800
 GCGGCATTAATAACAGACGGACCCGGTGGGGCAAAGCGAATGTATGTCCGCCGGCAAAC
 A A L I T D G P G G A K R M Y V G R Q N
 425

Fig. 2 (continued)

1810 1820 1830 1840 1850 1860
GCCGGTGAGACATGGCATGACATTACCGGAAACCGTTCGGAGCCGGTTGTCATCAATTCTG
A G E T W H D I T G N R S E P V V I N S
445
1870 1880 1890 1900 1910 1920
GAAGGCTGGGGAGAGTTTCACGTAAACGGCGGGTTCGGTTTCAATTTATGTTCAAAGATAG
E G W G E F H V N G G S V S I Y V Q R
465 483
1930 1940 1950 1960 1970 1980
AAGAGCAGAGAGGACGGATTTCCTGAAGGAAATCCGTTTTTTTATTTTGCCCGTCTTATA
1990 2000 2010 2020 2030 2040
AATTTCTTTGATTACATTTTATAATTAATTTTAACAAAGTGTTCATCAGCCCTCAGGAAGG
2050 2060 2070 2080 2090 2100
ACTTGCTGACAGTTTGAATCGCATAGGTAAGGCGGGGATGAAATGGCAACGTTATCTGAT
2110 2120 2130 2140
GTAGCAAAGAAAGCAAATGTGTCGAAAATGACGGTATCGCGGGTGATCA

Fig. 2 (continued)

```

      10      20      30      40      50      60
AATTCACCTCGAAAGCAAGCTGATAAACCGATACAATTAAAGGCTCCTTTTGGAGCCTTT

      70      80      90     100     110     120
TTTTTTGGAGATTTTCAACGTGAAAAAATTATTATTCGCAATTCCAAGCTAATTCACCTC

      130     140     150     160     170     180
GAAAGCAAGCTGATAAACCGATACAATTAAAGGCTCCTTTTGGAGCCTTTTTTTTGGAG

      190     200     210     220     230     240
ATTTTCAACGTGAAAAAATTATTATTCGCAATTCCAAGCTCTGCCTCGCGCGTTTCGGTG

      250     260     270     280     290     300
ATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGTCTGTAAG

      310     320     330     340     350     360
CGGATGCAGATCACGCGCCCTGTAGCGGCGCATTAAAGCGCGGCGGGTGTGGTGGTTACGC

      370     380     390     400     410     420
GCAGCGTGACCGCTACACTTGCCAGCGCCCTAGCGCCCGCTCCTTTTCGCTTCTTCCCTT

      430     440     450     460     470     480
CCTTTCTCGCCACGTTTCGCGGGCTTTCCCGTCAAGCTCTAAATCGGGGGCTCCCTTTAG

      490     500     510     520     530     540
GGTTCCGATTTAGTGCTTTACGGCACCTCGACCCCAAAAACTTGATTAGGGTGATGGTT

      550     560     570     580     590     600
CACGTAGTGGGCCATCGCCCTGATAGACGGTTTTTTCGCCCTTTGACGTTGGAGTCCACGT

      610     620     630     640     650     660
TCTTTAATAGTGGACTCTTGTTCCAAACTGGAACAACACTCAACCCTATCTCGGTCTATT

      670     680     690     700     710     720
CTTTTGATTTATAAGGGATTTTGCCGATTTTCGGCCTATTGGTTAAAAAATGAGCTGATTT

      730     740     750     760     770     780
AACAAAAATTTAACGCGAATTTTAACAAAATATTAACGTTTACAATTTGATCTGCGCTCG

      790     800     810     820     830     840
GTCGTTTCGGCTGCGGCGAGCGGTATCAGCTCACTCAAAGGCGGTAATACGGTTATCCACA

      850     860     870     880     890     900
GAATCAGGGGATAACGCAGGAAAGAACATGTGAGCAAAAGGCCAGCAAAAGGCCAGGAAC

      910     920     930     940     950     960
CGTAAAAAAGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCATCAC

      970     980     990    1000    1010    1020
AAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCAGGCG

      1030    1040    1050    1060    1070    1080
TTTCCCCCTGGAAGCTCCCTCGTGGCTCTCCTGTTCCGACCCTGCCGCTTACCGGATAC

      1090    1100    1110    1120    1130    1140
CTGTCCGCGCTTCTCCCTTCGGGAAGCGTGGCGCTTCTCAATGCTCACGCTGTAGGTAT

```

Figure 3

```

1150      1160      1170      1180      1190      1200
CTCAGTTCGGTGTAGGTCGTTTCGCTCCAAGCTGGGCTGTCTGCACGAACCCCCCGTTCAG

1210      1220      1230      1240      1250      1260
CCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACACGAC

1270      1280      1290      1300      1310      1320
TTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGGCGGT

1330      1340      1350      1360      1370      1380
GCTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATTTGGT

1390      1400      1410      1420      1430      1440
ATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGC

1450      1460      1470      1480      1490      1500
AAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTTGAAGCAGCAGATTACGCGCAGA

1510      1520      1530      1540      1550      1560
AAAAAAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAGTGGAAC

1570      1580      1590      1600      1610      1620
GAAAACTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTAGATC

1630      1640      1650      1660      1670      1680
CTTTTAAATTAAAAATGAAGTTTTTAAATCAATCTAAAGTATATATGAGTAACTTGGTCT

1690      1700      1710      1720      1730      1740
GACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGTCTATTTTCGTTCA

1750      1760      1770      1780      1790      1800
TCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACCATCT

1810      1820      1830      1840      1850      1860
GGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATCAGCA

1870      1880      1890      1900      1910      1920
ATAAACCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCCTGCAACTTTATCCGCCTCC

1930      1940      1950      1960      1970      1980
ATCCAGTCTATTAATTGTTGCCGGGAAGCTAGAGTAAGTAGTTCCGCCAGTTAATAGTTTG

1990      2000      2010      2020      2030      2040
CGCAACGTTGTTGCCATTGCTGCAGGCATCGTGGTGTACGCTCGTCGTTTGGTATGGCT

2050      2060      2070      2080      2090      2100
TCATTCAGCTCCGGTTCCTCAACGATCAAGGCCAGTTACATGATCCCCATGTTGTGCAAA

2110      2120      2130      2140      2150      2160
AAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAGTAAGTTGGCCGCGAGTGTTA

2170      2180      2190      2200      2210      2220
TCACTCATGGTTATGGCAGCACTGCATAATTCTCTTACTGTCATGCCATCCGTAAGATGC

2230      2240      2250      2260      2270      2280

```

Fig. 3 (continued)

TTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGGGGCGACCG
 2290 2300 2310 2320 2330 2340
 AGTTGCTCTTGGCCGGCGTCAACACGGGATAATACCGCCGCACATAGCAGAACTTTAAAA
 2350 2360 2370 2380 2390 2400
 GTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACTCTCAAGGATCTTACCGCTGTTG
 2410 2420 2430 2440 2450 2460
 AGATCCAGTTTCGATGTAACCCACTCGTGCACCCAACTGATCTTCAGCATCTTTTACTTTC
 2470 2480 2490 2500 2510 2520
 ACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAATAAGG
 2530 2540 2550 2560 2570 2580
 GCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCAGACAG
 2590 2600 2610 2620 2630 2640
 TTTTATTGTTTCATGATGATATATTTTTATCTTGTGCAATGTAAACATCAGAGATTTTGAGA
 2650 2660 2670 2680 2690 2700
 CACAACGTGGCTTTTGTGAATAAATCGAACTTTTGCTGAGTTGACTCCCCGCGCGCGATG
 2710 2720 2730 2740 2750 2760
 GGTCGAATTTGCTTTTCGAAAAAAGCCCGCTCATTAGGCGGGCTAAAAAAGCCCGCT
 2770 2780 2790 2800 2810 2820
 CATTAGGCGGGCTCGAATTTCTGCCATTTCATCCGCTTATTATCACTTATTCAGGCGTAGC
 2830 2840 2850 2860 2870 2880
 AACCAGGCGTTTAAGGGCACCAATAACTGCCCTTAAAAAATTACGCCCCGCCCTGCCACT
 2890 2900 2910 2920 2930 2940
 CATCGCAGTACTGTTGTAATTCATTAAGCATTCTGCCGACATGGAAGCCATCACAGACGG
 2950 2960 2970 2980 2990 3000
 CATGATGAACCTGAATCGCCAGCGGCATCAGCACCTTGTGCGCTTGGGTATAATATTTC
 3010 3020 3030 3040 3050 3060
 CCATAGTGAAAACGGGGGCGAAGAAGTTGTCCATATTCGCCACGTTTAAATCAAACTGG
 3070 3080 3090 3100 3110 3120
 TGAAACTCACCCAGGGATTGGCTGAGACGAAAAACATATTCTCAATAAACCTTTAGGGA
 3130 3140 3150 3160 3170 3180
 AATAGGCCAGGTTTTACCGTAACACGCCACATCTTGCGAATATATGTGTAGAACTGCC
 3190 3200 3210 3220 3230 3240
 GGAAATCGTCTGTTTCACTCCAGAGCGATGAAAACGTTTCAGTTTGGTTCATGGAAAA
 3250 3260 3270 3280 3290 3300
 CCGGTGTAACAAGGGTGAACACTATCCCATATCACCAGCTCACCGTCTTTCATTGCCATAC
 3310 3320 3330 3340 3350 3360
 GAAATTCGGGATGAGCATTTCATCAGGCGGGCAAGAATGTGAATAAAGGCCGATAAACT

Fig. 3 (continued)

3370 3380 3390 3400 3410 3420
 TGTGCTTATTTTCTTTACGGTCTTTAAAAAGGCCGTAATATCCAGCTAAACGGTCTGGT
 3430 3440 3450 3460 3470 3480
 TATAGGTACATTGAGCAACTGACTGAAATGCCTCAAAATGTTCTTTACGATGCCATTGGG
 3490 3500 3510 3520 3530 3540
 ATATATCAACGGTGGTATATCCAGTGATTTTTTTCTCCATTTTAGCTTCCTTAGCTCCTG
 3550 3560 3570 3580 3590 3600
 AAAATCTCGATAACTCAAAAAATACGCCCGGTAGTGATCTTATTTTCATTATGGTGAAAGT
 3610 3620 3630 3640 3650 3660
 TGGAACCTCTTACGTGCCGATCAACGTCTCATTTTCGCCAAAAGTTGGCCCAGGGCTTCC
 3670 3680 3690 3700 3710 3720
 CGGTATCAACAGGGACACCAGGATTTATTTATTCTGCGAAGTGATCTTCCGTCACAGGTA
 3730 3740 3750 3760 3770 3780
 TTTATTCGAAGACGAAAGGGCATCGCGCGCGGGAATTCGAGCTCGAGCTTACTCCCCAT
 3790 3800 3810 3820 3830 3840
 CCCCCTGTTGACAATTAATCATCGGCTCGTATAATGTGTGGAATTGTGAGCGGATAACAA
 3850 3860 3870 3880 3890 3900
 TTTACACAGGAAACAGGATCCAAGGAGGTGATCTAGAGTCATGAAACAACAAAAACGGC
 M K Q Q K R
 3910 3920 3930 3940 3950 3960
 TTTACGCCCGATTGCTGACGCTGTTATTTGCGCTCATCTTCTTGCTGCCTCATTTCTGCAG
 L Y A R L L T L L F A L I F L L P H S A
 3970 3980 3990 4000 4010 4020
 CAGCGGCGGCAAAATCTTAATGGGACGCTGATGCAGTATTTTGAATGGTACATGCCCAATG
 A A A A N L N G T L M Q Y F E W Y M P N
 +1
 4030 4040 4050 4060 4070 4080
 ACGGCCAACATTGGAAGCGTTTGCAAAACGACTCGGCATATTTGGCTGAACACGGTATTA
 D G Q H W K R L Q N D S A Y L A E H G I
 18
 4090 4100 4110 4120 4130 4140
 CTGCCGTCTGGATTCCCCCGCATATAAGGGAACTAGTCAAGCGGATGTGGGCTACGGTG
 T A V W I P P A Y K G T S Q A D V G Y G
 38
 4150 4160 4170 4180 4190 4200
 CTTACGACCTTTATGATTTAGGGGAGTTTCATCAAAAAGGGACGGTTCCGACAAAGTACG
 A Y D L Y D L G E F H Q K G T V R T K Y
 58
 4210 4220 4230 4240 4250 4260
 GCACAAAAGGAGAGCTGCAATCTGCGATCAAAAAGTCTTCATTCCCGCGACATTAACGTTT
 G T K G E L Q S A I K S L H S R D I N V
 78
 4270 4280 4290 4300 4310 4320
 ACGGGGATGTGGTCATCAACCACAAAGGCGCGCTGATGCGACCGAAGATGTAACCGCGG
 Y G D V V I N H K G G A D A T E D V T A
 98

Fig. 3 (continued)

4330 4340 4350 4360 4370 4380
 TTGAAGTCGATCCCGCTGACCGCAACCGCGTAATTTTCAGGAGAACACCTAATTAAAGCCT
 V E V D P A D R N R V I S G E H L I K A
 118

4390 4400 4410 4420 4430 4440
 GGACACATTTTCATTTTCCGGGGCGCGGCAGCACATACAGCGATTTTAAATGGCATTGGT
 W T H F H F P G R G S T Y S D F K W H W
 138

4450 4460 4470 4480 4490 4500
 ACCATTTTGACGGAACCGATTGGGACGAGTCCCGAAAGCTGAACCGCATCTATAAGTTTC
 Y H F D G T D W D E S R K L N R I Y K F
 158

4510 4520 4530 4540 4550 4560
 AAGGAAAGGCTTGGGATTGGGAAGTTTCCAATGAAAACGGCAACTATGATTATTTGATGT
 Q G K A W D W E V S N E N G N Y D Y L M
 178

4570 4580 4590 4600 4610 4620
 ATGCCGACATCGATTATGACCATCCTGATGTTCGCAGCAGAAATTAAGAGATGGGGCACTT
 Y A D I D Y D H P D V A A E I K R W G T
 198

4630 4640 4650 4660 4670 4680
 GGTATGCCAATGAACTGCAATTGGACGGTTTTCCGTCTTGATGCTGTCAAACACATTAAAT
 W Y A N E L Q L D G F R L D A V K H I K
 218

4690 4700 4710 4720 4730 4740
 TTTCTTTTTTTCGGGATTGGGTTAATCATGTTCAGGGAAAAACGGGGAAGGAAATGTTTA
 F S F L R D W V N H V R E K T G K E M F
 238

4750 4760 4770 4780 4790 4800
 CGGTAGCTGAATATTGGCAGAATGACTTGGGCGCCCTGGAAAACTATTTGAACAAAACAA
 T V A E Y W Q N D L G A L E N Y L N K T
 258

4810 4820 4830 4840 4850 4860
 ATTTTAATCATTGAGTGTGACGTGCCGCTTCATTATCAGTTCCATGCTGCATCGACAC
 N F N H S V F D V P L H Y Q F H A A S T
 278

4870 4880 4890 4900 4910 4920
 AGGGAGGCGGCTATGATATGAGGAAATTGCTGAACGGTACGGTCGTTTCCAAGCATCCGT
 Q G G G Y D M R K L L N G T V V S K H P
 298

4930 4940 4950 4960 4970 4980
 TGAAATCGGTTACATTTGTCGATAACCATGATACACAGCCGGGGCAATCGCTTGAGTCGA
 L K S V T F V D N H D T Q P G Q S L E S
 318

4990 5000 5010 5020 5030 5040
 CTGTCCAAACATGGTTAAGCCGCTTGCTTACGCTTTTATTCTCACAAGGGAATCTGGAT
 T V Q T W F K P L A Y A F I L T R E S G
 338

5050 5060 5070 5080 5090 5100
 ACCCTCAGGTTTTCTACGGGGATATGTACGGGACGAAAGGAGACTCCCAGCGCGAAATTC
 Y P Q V F Y G D M Y G T K G D S Q R E I
 358

5110 5120 5130 5140 5150 5160
 CTGCCTTGAAACACAAAATTGAACCGATCTTAAAAGCGAGAAAACAGTATGCGTACGGAG
 P A L K H K I E P I L K A R K Q Y A Y G
 378

Fig. 2 (continued)

5170 5180 5190 5200 5210 5220
CACAGCATGATTATTTTCGACCACCATGACATTGTCGGCTGGACAAGGGAAGGCCGACAGCT
A Q H D Y F D H H D I V G W T R E G D S
398

5230 5240 5250 5260 5270 5280
CGGTTGCAAATTCAGGTTTGGCGGCATTAATAACAGACGGACCCGGTGGGGCAAAGCGAA
S V A N S G L A A L I T D G P G G A K R
418

5290 5300 5310 5320 5330 5340
TGTATGTCGGCCGGCAAACGCCGGTGAGACATGGCATGACATTACCGGAAACCGTTCGG
M Y V G R Q N A G E T W H D I T G N R S
438

5350 5360 5370 5380 5390 5400
AGCCGGTTGTCATCAATTCGGAAGGCTGGGGAGAGTTTCACGTAAACGGCGGGTTCGGTTT
E P V V I N S E G W G E F H V N G G S V
458

5410 5420 5430 5440 5450 5460
CAATTTATGTTCAAAGATAGGTGACCAGAGAGGACGGATTTCCTGAAGGAAATCCGTTTT
S I Y V Q R
478

5470 5480 5490 5500 5510 5520
TTTATTTTGCCCGTCTTATAAATTTCTTTGATTACATTTTATAATTAATTTTAACAAAGT

5530 5540 5550 5560 5570 5580
GTCATCAGCCCTCAGGAAGGACTTGCTGACAGTTTGAATCGCATAGGTAAGGCGGGGATG

5590 5600 5610 5620 5630 5640
AAATGGCAACGTTATCTGATGTAGCAAAGAAAGCAAATGTGTCGAAAATGACGGTATCGC

5650 5660 5670
GGGTGATCCTCTAGAAGAAGCTTGGTCTAGAGGTCGA

Fig. 3 (continued)